
COMPUTER SYSTEM
100
PROCESSOR
102
ARITHMETIC
UNIT
104
MEMORY 112
COMPUTATIONAL
CODE FOR
INTERVALS
114
INTERVAL VALUES
115
BRIDGE
106
PERIPHERAL BUS
110
STORAGE
DEVICE
108

FIG. 1


INTERVAL
VALUES
115
PROCESSOR
102
SOURCE CODE
FOR INTERVAL
COMPUTATIONS
202
COMPILER FOR
INTERVAL CODE
204
EXECUTABLE
CODE FOR
INTERVAL
COMPUTATIONS
206
ARITHMETIC
UNIT FOR
INTERVAL
COMPUTATIONS
104
RESULTS
212

FIG. 2


INTERVAL 322
FIRST FLOATING POINT NUMBER 324
SECOND FLOATING POINT NUMBER 326
ARITHMETIC UNIT FOR INTERVAL COMPUTATIONS 104
FIRST FLOATING POINT NUMBER 304
SECOND FLOATING POINT NUMBER 306
FIRST FLOATING POINT NUMBER 314
SECOND FLOATING POINT NUMBER 316
INTERVAL 302
INTERVAL 312

FIG. 3


START 400
RECEIVE REPRESENTATION OF INTERVAL 402
PERFORM ARITHMETIC OPERATION USING REPRESENTATION OF INTERVAL 404
END 406

FIG. 4

$$X \equiv \left[\underline{x}, \bar{x}\right] \equiv \left\{x \in \Re^* \middle| \underline{x} \le x \le \bar{x}\right\}$$

$$Y \equiv \left[\underline{y}, \bar{y}\right] \equiv \left\{y \in \Re^* \middle| \underline{y} \le y \le \bar{y}\right\}$$

(1) $$X + Y = \left[\downarrow \underline{x} + \underline{y}, \uparrow \bar{x} + \bar{y}\right]$$

(2) $$X - Y = \left[\downarrow \underline{x} - \bar{y}, \uparrow \bar{x} - \underline{y}\right]$$

(3) $$X \times Y = \left[\min\left(\downarrow \underline{x} \times \underline{y}, \underline{x} \times \bar{y}, \bar{x} \times \underline{y}, \bar{x} \times \bar{y}\right), \max\left(\uparrow \underline{x} \times \underline{y}, \underline{x} \times \bar{y}, \bar{x} \times \underline{y}, \bar{x} \times \bar{y}\right)\right]$$

(4) $$X / Y = \left[\min\left(\downarrow \underline{x} / \underline{y}, \underline{x} / \bar{y}, \bar{x} / \underline{y}, \bar{x} / \bar{y}\right), \max\left(\uparrow \underline{x} / \underline{y}, \underline{x} / \bar{y}, \bar{x} / \underline{y}, \bar{x} / \bar{y}\right)\right], \text{ if } 0 \notin Y$$

$$X / Y \subseteq \Re^*, \text{ if } 0 \in Y$$

FIG. 5


START
RECEIVE REPRESENTATION OF NONLINEAR EQUATION $f(\mathbf{x})=0$ 602
RECEIVE REPRESENTATION OF INITIAL BOX $\mathbf{X}$ WITH X_j A COMPONENT OF $\mathbf{X}$ 604
SYMBOLICALLY MANIPULATE $f(\mathbf{x})=0$ TO SOLVE FOR A TERM $g(x_j)=h(\mathbf{x})$, WHEREIN $g(x_j)$ CAN BE ANALYTICALLY INVERTED 606
SUBSTITUTE $\mathbf{X}$ INTO $g(X'_j)=h(\mathbf{X})$ 608
SOLVE FOR $X'_j = g^{-1}[h(\mathbf{X})]$ 610
INTERSECT X'_j WITH X_j TO PRODUCE NEW INTERVAL X_j^+ 612
$X_j = X_j^+$ 614
WAS THE BOX SUFFICIENTLY REDUCED? 616
YES
NO
IF X_j^+ IS EMPTY
END
REPEAT FOR OTHER TERMS AND OTHER VALUES OF j

FIG. 6


START
FOR EACH BOX IN LIST L_1, APPLY TERM CONSISTENCY TO EACH OF THE INEQUALITY CONSTRAINTS $p_i(\mathbf{x}) \le 0$ (i=1, ..., m).
701
IF f_bar <+∞, THEN FOR EACH BOX IN L_1, APPLY TERM CONSISTENCY TO THE INEQUALITY $f \le f\_bar$.
702
IF L_1 IS EMPTY, GO TO STEP 742. OTHERWISE, SELECT THE BOX IN L_1 FOR WHICH THE LOWER BOUND OF $f(\mathbf{X})$ IS SMALLEST. FOR LATER REFERENCE, DENOTE THIS BOX BY $\mathbf{X}^{(1)}$. DELETE $\mathbf{X}$ FROM L_1.
703
APPLY TERM CONSISTENCY OVER $\mathbf{X}$ TO EACH CONSTRAINT INEQUALITY. IF $\mathbf{X}$ IS DELETED, GO TO STEP 703.
704
COMPUTE AN APPROXIMATION $\mathbf{x}$ FOR THE CENTER $m(\mathbf{X})$ OF $\mathbf{X}$. IF $f(\mathbf{x}) > f\_bar$, GO TO STEP 708.
705
FOR FUTURE REFERENCE, DENOTE THE BOX $\mathbf{X}$ by $\mathbf{X}^{(2)}$. DO A CONSTRAINED LINE SEARCH TO TRY TO REDUCE f_bar.
706
IF f_bar WAS NOT REDUCED IN STEP 706, GO TO STEP 709.
707
APPLY TERM CONSISTENCY TO THE INEQUALITY $f(\mathbf{x}) \le f\_bar$ OVER THE CURRENT BOX $\mathbf{X}$. IF $\mathbf{X}$ IS DELETED, GO TO STEP 703.
708
IF $w(\mathbf{X}) < \varepsilon_X$ AND $w[f(\mathbf{X})] < \varepsilon_F$, PUT $\mathbf{X}$ IN LIST L_2. OTHERWISE, IF $\mathbf{X}$ IS SUFFICIENTLY REDUCED RELATIVE TO THE BOX $\mathbf{X}^{(1)}$, PUT $\mathbf{X}$ IN L_1 AND GO TO STEP 703.
709
APPLY BOX CONSISTENCY TO EACH INEQUALITY CONSTRAINT. IF f_bar < +∞, APPLY BOX CONSISTENCY TO THE INEQUALITY $f(\mathbf{x}) \le f\_bar$. IF $\mathbf{X}$ IS DELETED, GO TO STEP 703.
710
IF THE UPPER BOUND OF $p_i(\mathbf{X}) \ge 0$ FOR ANY i=1, ..., n, GO TO STEP 726.
711
APPLY TERM CONSISTENCY TO g_i=0 FOR i=1, ..., n. IF THE RESULT FOR ANY i=1, ..., n IS EMPTY, GO TO STEP 703.
712
A

FIG. 7A


A
APPLY TERM CONSISTENCY TO THE RELATION $H_{ii}(\mathbf{x}) \geq 0$ FOR $i=1, ..., n$. IF THE RESULT IS EMPTY, GO TO STEP 703.
713
REPEAT STEP 709.
714
APPLY BOX CONSISTENCY TO $g_i=0$ FOR $i=1, ..., n$. IF THE RESULT IS EMPTY, GO TO STEP 703.
715
APPLY BOX CONSISTENCY TO $H_{ii}(\mathbf{x}) \geq 0$ FOR $i=1, ..., n$. IF THE RESULT IS EMPTY, GO TO STEP 703.
716
REPEAT STEP 709.
717
USE A CRITERION $w(\mathbf{X}) > (w_l+w_R)/2$ TO DECIDE IF A NEWTON STEP SHOULD BE APPLIED TO SOLVE $\mathbf{g}=\mathbf{0}$. IF NOT, GO TO STEP 726.
718
GENERATE THE INTERVAL JACOBIAN $\mathbf{J}(\mathbf{x},\mathbf{X})$ OF THE GRADIENT $\mathbf{g}$ AND COMPUTE THE APPROXIMATE INVERSE $\mathbf{B}$ OF THE CENTER OF $\mathbf{J}(\mathbf{x},\mathbf{X})$. APPLY ONE STEP OF AN INTERVAL NEWTON METHOD TO SOLVE $\mathbf{g}=\mathbf{0}$. IF THE RESULT IS EMPTY, GO TO STEP 703.
719
REPEAT STEP 709.
720
OBTAIN $\mathbf{Bg}$ IN ANALYTIC FORM. APPLY TERM CONSISTENCY TO SOLVE THE i^{th} EQUATION OF $\mathbf{Bg}=0$ FOR THE i^{th} VARIABLE x_i FOR $i=1, ..., n$. IF THE RESULT IS EMPTY, GO TO STEP 703.
721
REPEAT STEP 709.
722
USE BOX CONSISTENCY TO SOLVE THE i^{th} EQUATION OF $\mathbf{Bg}$ FOR THE i^{th} VARIABLE FOR $i=1, ..., n$. IF THE RESULT IS EMPTY, GO TO STEP 703.
723
REPEAT STEP 709.
724
B

FIG. 7B


B
USE THE MATRIX B IN A NEWTON STEP TO TRY TO REDUCE f_bar.
725
COMPUTE AN APPROXIMATION x FOR THE CENTER m(X) OF X. IF f(x)>f_bar, GO TO STEP 703.
726
SKIP THIS STEP AND GO TO STEP 732 IF X=X(2) IS THE SAME BOX FOR WHICH A LINE SEARCH WAS DONE IN STEP 706. OTHERWISE, DO A LINE SEARCH TO TRY TO REDUCE f_bar. IF f_bar IS NOT REDUCED, GO TO STEP 732.
727
FOR FUTURE REFERENCE DENOTE X(3) = X. USE A LINEARIZATION TEST TO DECIDE WHETHER TO LINEARIZE AND "SOLVE" THE INEQUALITY f(x) ≤ f_bar. IF THE CRITERION IS NOT SATISFIED, GO TO STEP 732.
728
USE A LINEAR METHOD TO TRY TO REDUCE X USING THE INEQUALITY f(x) ≤ f_bar. IF X IS DELETED, GO TO STEP 703. OTHERWISE, IF THIS APPLICATION OF THE LINEAR METHOD DOES NOT SUFFICIENTLY REDUCE BOX X(3) GO TO STEP 731.
729
USE A QUADRATIC METHOD TO TRY TO REDUCE X USING THE INEQUALITY f(x) ≤ f_bar. IF X IS DELETED, GO TO STEP 703.
730
REPEAT STEP 709.
731
USE A LINEARIZATION TEST TO DECIDE WHETHER TO LINEARIZE AND "SOLVE" THE INEQUALITY CONSTRAINTS. IF THE PROCEDURE INDICATES THAT THE LINEARIZATION SHOULD NOT BE DONE, GO TO STEP 739.
732
SELECT THE INEQUALITY CONSTRAINTS TO BE SOLVED IN LINEARIZED FORM, AND POSSIBLY ADD TO THIS SET THE INEQUALITY f(x) ≤ f_bar. IF NO INEQUALITIES ARE SELECTED, GO TO STEP 739. OTHERWISE, LINEARIZE THE RESULTING SET OF INEQUALITIES, AND SOLVE THE RESULTING SET OF LINEAR INEQUALITIES. IF THE SOLUTION SET IS EMPTY, GO TO STEP 703.
733
REPEAT STEP 709.
734
C

FIG. 7C


C
USE PRECONDITIONING MATRIX B TO ANALYTICALLY PRECONDITION THE SET OF INEQUALITIES THAT WERE SELECTED FOR USE IN STEP 733. USE TERM CONSISTENCY TO SOLVE EACH PRECONDITIONED INEQUALITY.
735
REPEAT STEP 709.
736
USE BOX CONSISTENCY TO SOLVE THE SAME INEQUALITIES FOR THE SAME VARIABLES AS IN STEP 735.
737
REPEAT STEP 709.
738
USE A LINEARIZATION TEST TO DECIDE WHETHER TO SOLVE THE JOHN CONDITIONS. IF NOT, GO TO STEP 742.
739
MODIFY THE JOHN CONDITIONS BY OMITTING THOSE CONSTRAINTS p_i FOR WHICH $p_i(\mathbf{X}) < 0$ (SINCE THEY ARE NOT ACTIVE IN X). APPLY ONE PASS OF THE INTERVAL NEWTON METHOD TO THE (MODIFIED) JOHN CONDITIONS. IF THE RESULT IS EMPTY, GO TO STEP 703.
740
REPEAT STEP 709.
741
IN VARIOUS PREVIOUS STEPS, GAPS MAY HAVE BEEN FORMED IN COMPONENTS OF X. IF SO, MERGE ANY OVERLAPPING GAPS. SPLIT X, AND PLACE THE RESULTING SUBBOXES IN L_1 AND GO TO STEP 703.
742
IF $f\_bar < +\infty$, APPLY TERM CONSISTENCY TO $f(\mathbf{x}) \le f\_bar$ FOR EACH BOX IN THE LIST L_2. DENOTE THOSE THAT REMAIN BY $\mathbf{X}^{(1)}, \ldots, \mathbf{X}^{(s)}$ WHERE s IS THE NUMBER OF BOXES REMAINING.. DETERMINE
$\underline{F} = \min_{1 \le i \le s} \underline{f}(X^{(i)})$ and $\overline{F} = \max_{1 \le i \le s} \overline{f}(X^{(i)})$
743
TERMINATE.
744

FIG. 7D